




# California River Indices

## California River Indices

Hydrology: California's water development has generally been dictated by extremes of droughts and floods. The six-year drought of 1929-34 established the criteria commonly used to plan storage capacity or water yield of large Northern California reservoirs. The influence of climatic variability on California's water supplies is much less predictable than the influences of geographic and seasonal variability, as evidenced by the recent historical records of precipitation and runoff. For example, the state's average annual runoff includes the all-time low of 15 million acre-feet in 1977 and the all-time high of over 135 million acre-feet in 1983. Floods and droughts occur often, sometimes in the same year. The January 1997 flood was followed by a record-setting dry period from February through June, and the flooding of 1986 was followed by six years of drought (1987-92).


Figures showing the estimated annual unimpaired runoff of the Sacramento and San Joaquin River basins illustrate climatic variability. Because these basins provide much of the state's water supply, their hydrology is often used as indices of water year classification systems.

Unimpaired runoff represents the natural water production of a river basin, unaltered by upstream diversions, storage, and export of water to or import of water from other basins.



Unimpaired runoff represents the natural water production of a river basin, unaltered by upstream diversions, storage, export of water to or import of water from other basins.

Sacramento River Runoff is the sum (in million acre-feet) of Sacramento River at Bend Bridge, Feather River inflow to Lake Oroville, Yuba River at Smartville, and American River inflow to Folsom Lake. The water year sum is also known as the Sacramento River Index, and was previously referred to as the "4 River Index" or "4 Basin Index". It was previously used to determine year-type classifications under SWRCB Decision 1485.



San Joaquin River Runoff is the sum of Stanislaus River inflow to New Melones Lake, Tuolumne River inflow to New Don Pedro Reservoir, Merced River inflow to Lake McClure, and San Joaquin River inflow to Millerton Lake (all in million acre-feet).

**Water Year Classification:** Water-year classification systems provide a means to assess the amount of water originating in a basin. Because water-year classification systems are useful in water planning and management, they have been developed for several hydrologic basins in California. The Sacramento Valley 40-30-30 Index and the San Joaquin Valley 60-20-20 Index were developed by the State Water Resources Control Board (SWRCB) for the Sacramento and San Joaquin River hydrologic basins as part of SWRCB's Bay-Delta regulatory activities. Both systems define one "wet" year classification, two "normal" classifications (above and below normal), and two "dry" classifications (dry and critical), for a total of five water year types.

Sacramento Valley Water Year Index =  $(0.4) \times \text{Current Apr-Jul runoff forecast (in million acre-feet)} + (0.3) \times \text{Current Oct-Mar runoff (in million acre-feet)} + (0.3) \times \text{Previous Water Year's Index (if the Previous Water Year's Index exceeds 10.0, then 10.0 is used)}$ .

This index, originally specified in the 1995 SWRCB Water Quality Control Plan, is used to determine the Sacramento Valley water-year type as implemented in SWRCB D-1641. Year types are set by first of month forecasts beginning in February. Final determination is based on the May 1, 50 percent exceedence forecast.

